SPECIES SPECIFIC AGAINST: PEST BIRDS in commercial, industrial, rural and residential situations. Do not use when gulls are present.

PREPARING SURFACES FOR TREATMENT

Ensure appropriate PPE is used when preparing the site, removal of all nests and other debris. For effective installation of Ornaway Bird Gel all surfaces within the installation area must be thoroughly cleaned and free from faecal and organic matter. In some cases where birds have been habitually nesting in an area for several years (high pressure sites), these birds will need to be either trapped or despatched before installation (Check national legislation and guidelines for humane despatching) or Ornaway should be installed in combination with appropriate netting and spikes as part of an integrated plan to keep birds away from a specific area.

Enhanced efficacy will be gained by treating the cleaned areas with a fine mist spray of disinfectant (e.g. Guanaway). Do not over-wet the area as this will lengthen drying time and inhibit adhesion of silicone. Ornaway Bird Gel will only be completely effective if the site is properly cleaned before installation.

APPLICATION

Following surface preparation of the treatment area, apply dishes to surfaces according to the spacing and application rate guidelines in a grid fashion. Dishes can be stuck to the surface with an appropriate adhesive.

STORAGE

Cool dry conditions. Keep out of reach of children.

DISPOSAL

Product contains food and pharmaceutical grade materials and can be disposed of appropriately on site or at an approved waste disposal facility. Never dispose unused product down any indoor or outdoor drains.

PRE-FILLED

Ornaway ®
Bird Gel | Visual Bird Repellent

SPACING & APPLICATION RATES

High pressure site
4-6 inches (10-15cm)

Medium pressure site
8-10 inches (20-25cm)

Low pressure site
12-20 inches (30-50cm)

Barrettine *Environmental Health*

R2